

PARAMETER ESTIMATION OF GAMMA-PARETO DISTRIBUTION VIA BAYESIAN APPROACH

Arun Kumar Rao & Himanshu Pandey

Department of Mathematics & Statistics, DDU Gorakhpur University, Gorakhpur, INDIA
E-mail: himanshu_pandey62@yahoo.com

ARTICLE INFO

Received: 28 February 2022
Revised: 30 April 2022
Accepted: 18 May 2022
Online: 23 June 2022

To cite this paper:
Arun Kumar Rao &
Himanshu Pandey (2022).
Parameter Estimation of
Gamma-Pareto
Distribution Via Bayesian
Approach. *International
Journal of Mathematics,
Statistics and Operations
Research*. 2(1): pp. 61-70.
[https://DOI: 10.47509 /
IJMSOR.2022.v02i01.05](https://DOI: 10.47509/IJMSOR.2022.v02i01.05)

ABSTRACT

In this paper, Gamma-Pareto distribution is considered for Bayesian analysis. The expressions for Bayes estimators of the parameter have been derived under squared error, precautionary, entropy, K-loss, and Al-Bayyati's loss functions by using quasi and inverted gamma priors.

Keywords: Bayesian method, Gamma-Pareto distribution, quasi and inverted gamma priors, squared error, precautionary, entropy, K-loss, and Al-Bayyati's loss functions.

1. INTRODUCTION

The Gamma-Pareto distribution was proposed by Alzaatreh et al. [1]. They obtained the various properties of this distribution and the method of maximum likelihood was proposed for estimating the parameters. The probability density function of Gamma-Pareto distribution is given by

$$f(x; \theta) = \frac{1}{\Gamma(\lambda)} \theta^{-\lambda} x^{-1} \left(\frac{c}{x}\right)^{(1/\theta)} \left[\log\left(\frac{x}{c}\right)\right]^{\lambda-1}; x > 0. \quad (1)$$

The joint density function or likelihood function of (1) is given by

$$f(\underline{x}; \theta) = \left(\frac{1}{\Gamma(\lambda)}\right)^n \left(\prod_{i=1}^n x_i^{-1} \left[\log\left(x_i/c\right)\right]^{\lambda-1}\right) \exp\left[-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right]. \quad (2)$$

The log likelihood function is given by

$$\log f(x; \theta) = n \log \left(\frac{1}{\Gamma(\lambda)} \right) - n\lambda \log \theta + \log \left(\prod_{i=1}^n x_i^{-1} [\log(x_i/c)]^{\lambda-1} \right) - \frac{1}{\theta} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right). \quad (3)$$

Differentiating (3) with respect to θ and equating to zero, we get the maximum likelihood estimator of θ which is given as

$$\hat{\theta} = \frac{1}{n\lambda} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right). \quad (4)$$

2. BAYESIAN METHOD OF ESTIMATION

The Bayesian inference procedures have been developed generally under squared error loss function

$$L(\hat{\theta}, \theta) = (\hat{\theta} - \theta)^2. \quad (5)$$

The Bayes estimator under the above loss function, say, $\hat{\theta}_s$ is the posterior mean, i.e,

$$\hat{\theta}_s = E(\theta). \quad (6)$$

Zellner [2], Basu and Ebrahimi [3] have recognized that the inappropriateness of using symmetric loss function. Norstrom [4] introduced precautionary loss function is given as

$$L(\hat{\theta}, \theta) = \frac{(\hat{\theta} - \theta)^2}{\hat{\theta}}. \quad (7)$$

The Bayes estimator under this loss function is denoted by $\hat{\theta}_p$ and is obtained as

$$\hat{\theta}_p = [E(\theta^2)]^{1/2}. \quad (8)$$

Calabria and Pulcini [5] points out that a useful asymmetric loss function is the entropy loss

$$L(\delta) \propto [\delta^p - p \log_e(\delta) - 1]$$

where $\delta = \frac{\hat{\theta}}{\theta}$, and whose minimum occurs at $\hat{\theta} = \theta$. Also, the loss function

$L(\delta)$ has been used in Dey et al. [6] and Dey and Liu [7], in the original form having $p = 1$. Thus can written be as

$$L(\delta) = b[\delta - \log_e(\delta) - 1]; \quad b > 0. \quad (9)$$

The Bayes estimator under entropy loss function is denoted by $\hat{\theta}_E$ and is obtained by solving the following equation

$$\hat{\theta}_E = \left[E\left(\frac{1}{\theta}\right) \right]^{-1}. \quad (10)$$

Wasan [8] proposed the K-loss function which is given as

$$L(\hat{\theta}, \theta) = \frac{(\hat{\theta} - \theta)^2}{\hat{\theta}\theta}. \quad (11)$$

Under K-loss function the Bayes estimator of θ is denoted by $\hat{\theta}_K$ and is obtained as

$$\hat{\theta}_K = \left[\frac{E(\theta)}{E(1/\theta)} \right]^{\frac{1}{2}}. \quad (12)$$

Al-Bayyati [9] introduced a new loss function which is given as

$$L(\hat{\theta}, \theta) = \theta^c (\hat{\theta} - \theta)^2. \quad (13)$$

Under Al-Bayyati's loss function the Bayes estimator of θ is denoted by $\hat{\theta}_{Al}$ and is obtained as

$$\hat{\theta}_{AI} = \frac{E(\theta^{c+1})}{E(\theta^c)}. \quad (14)$$

Let us consider two prior distributions of θ to obtain the Bayes estimators.

- (i) Quasi-prior: For the situation where we have no prior information about the parameter θ , we may use the quasi density as given by

$$g_1(\theta) = \frac{1}{\theta^d}; \theta > 0, d \geq 0, \quad (15)$$

where $d = 0$ leads to a diffuse prior and $d = 1$, a non-informative prior.

- (ii) Inverted gamma prior: Generally, the inverted gamma density is used as prior distribution of the parameter θ given by

$$g_2(\theta) = \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{-(\alpha+1)} e^{-\beta/\theta}; \theta > 0. \quad (16)$$

3. POSTERIOR DENSITY UNDER $g_1(\theta)$

The posterior density of θ under $g_1(\theta)$, on using (2), is given by

$$\begin{aligned} f(\theta/\underline{x}) &= \frac{\left(\frac{1}{\Gamma(\lambda)}\right)^n \theta^{-n\lambda} \left(\prod_{i=1}^n x_i^{-1} [\log(x_i/c)]^{\lambda-1}\right) \exp\left[-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right] \theta^{-d}}{\int_0^\infty \left(\frac{1}{\Gamma(\lambda)}\right)^n \theta^{-n\lambda} \left(\prod_{i=1}^n x_i^{-1} [\log(x_i/c)]^{\lambda-1}\right) \exp\left[-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right] \theta^{-d} d\theta} \\ &= \frac{\theta^{-(n\lambda+d)} e^{-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)}}{\int_0^\infty \theta^{-(n\lambda+d)} e^{-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)} d\theta} \\ &= \frac{\left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+d-1}}{\Gamma(n\lambda+d-1)} \theta^{-(n\lambda+d)} e^{-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)} \end{aligned} \quad (17)$$

Theorem 1. On using (17), we have

$$E(\theta^c) = \frac{\Gamma(n\lambda + d - c - 1)}{\Gamma(n\lambda + d - 1)} \left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^c \quad (18)$$

Proof. By definition,

$$\begin{aligned} E(\theta^c) &= \int \theta^c f(\theta/\underline{x}) d\theta \\ &= \frac{\left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^{n\lambda+d-1}}{\Gamma(n\lambda + d - 1)} \int_0^\infty \theta^{-(n\lambda+d-c)} e^{-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)} d\theta \\ &= \frac{\left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^{n+d-1}}{\Gamma(n\lambda + d - 1)} \frac{\Gamma(n\lambda + d - c - 1)}{\left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^{n\lambda+d-c-1}} \\ &= \frac{\Gamma(n\lambda + d - c - 1)}{\Gamma(n\lambda + d - 1)} \left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^c. \end{aligned}$$

From equation (18), for $c = 1$, we have

$$E(\theta) = \frac{1}{(n\lambda + d - 2)} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \quad (19)$$

From equation (18), for $c = 2$, we have

$$E(\theta^2) = \frac{1}{(n\lambda + d - 2)(n\lambda + d - 3)} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \quad (20)$$

From equation (18), for $c = -1$, we have

$$E\left(\frac{1}{\theta}\right) = (n\lambda + d - 1) \left(\sum_{i=1}^n \log\left(\frac{x_i}{c}\right) \right)^{-1} \quad (21)$$

From equation (18), for $c = c + 1$, we have

$$E(\theta^{c+1}) = \frac{\Gamma(n\lambda + d - c - 2)}{\Gamma(n\lambda + d - 1)} \left(\sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \right)^{c+1} \quad (22)$$

4. BAYES ESTIMATORS UNDER $g_1(\theta)$

From equation (6), on using (19), the Bayes estimator of θ under squared error loss function is given by

$$\hat{\theta}_S = \frac{1}{(n\lambda + d - 2)} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \quad (23)$$

From equation (8), on using (20), the Bayes estimator of θ under precautionary loss function is given by

$$\hat{\theta}_P = \left[(n\lambda + d - 2)(n\lambda + d - 3) \right]^{-\frac{1}{2}} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \quad (24)$$

From equation (10), on using (21), the Bayes estimator of θ under entropy loss function is given by

$$\hat{\theta}_E = \frac{1}{(n\lambda + d - 1)} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right). \quad (25)$$

From equation (12), on using (19) and (21), the Bayes estimator of θ under K-loss function is given by

$$\hat{\theta}_K = \left[(n\lambda + d - 2)(n\lambda + d - 1) \right]^{-\frac{1}{2}} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \quad (26)$$

From equation (14), on using (18) and (22), the Bayes estimator of θ under Al-Bayyati's loss function is given by

$$\hat{\theta}_{Al} = \frac{1}{(n\lambda + d - c - 2)} \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \quad (27)$$

5. POSTERIOR DENSITY UNDER $g_2(\theta)$

Under $g_2(\theta)$, the posterior density of θ , using equation (2), is obtained as

$$\begin{aligned}
f(\theta/\underline{x}) &= \frac{\left(\frac{1}{\Gamma(\lambda)}\right)^n \theta^{-n\lambda} \left(\prod_{i=1}^n x_i^{-1} [\log(x_i/c)]^{\lambda-1}\right) \exp\left[-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right] \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{-(\alpha+1)} e^{-\beta/\theta}}{\int_0^\infty \left(\frac{1}{\Gamma(\lambda)}\right)^n \theta^{-n\lambda} \left(\prod_{i=1}^n x_i^{-1} [\log(x_i/c)]^{\lambda-1}\right) \exp\left[-\frac{1}{\theta} \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right] \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{-(\alpha+1)} e^{-\beta/\theta} d\theta} \\
&= \frac{\theta^{-(n\lambda+\alpha+1)} e^{-\frac{1}{\theta}\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)}}{\int_0^\infty \theta^{-(n\lambda+\alpha+1)} e^{-\frac{1}{\theta}\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)} d\theta} \\
&= \frac{\theta^{-(n\lambda+\alpha+1)} e^{-\frac{1}{\theta}\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)}}{\Gamma(n\lambda+\alpha) \left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+\alpha}} \\
&= \frac{\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+\alpha}}{\Gamma(n\lambda+\alpha)} \theta^{-(n\lambda+\alpha+1)} e^{-\frac{1}{\theta}\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)}. \tag{28}
\end{aligned}$$

Theorem 2. On using (28), we have

$$E(\theta^c) = \frac{\Gamma(n\lambda+\alpha-c)}{\Gamma(n\lambda+\alpha)} \left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^c \tag{29}$$

Proof. By definition,

$$E(\theta^c) = \int \theta^c f(\theta/\underline{x}) d\theta$$

$$= \frac{\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+\alpha}}{\Gamma(n\lambda+\alpha)} \int_0^\infty \theta^{-(n\lambda+\alpha+1-c)} e^{-\frac{1}{\theta}\left(\beta+\sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)} d\theta$$

$$\begin{aligned}
&= \frac{\left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+\alpha}}{\Gamma(n\lambda+\alpha)} \frac{\Gamma(n\lambda+\alpha-c)}{\left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{n\lambda+\alpha-c}} \\
&= \frac{\Gamma(n\lambda+\alpha-c)}{\Gamma(n\lambda+\alpha)} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^c
\end{aligned}$$

From equation (29), for $c = 1$, we have

$$E(\theta) = \frac{1}{(n\lambda+\alpha-1)} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right) \quad (30)$$

From equation (29), for $c = 2$, we have

$$E(\theta^2) = [(n\lambda+\alpha-1)(n\lambda+\alpha-2)]^{-1} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^2 \quad (31)$$

From equation (29), for $c = -1$, we have

$$E\left(\frac{1}{\theta}\right) = (n\lambda+\alpha) \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{-1} \quad (32)$$

From equation (29), for $c = c+1$, we have

$$E(\theta^{c+1}) = \frac{\Gamma(n\lambda+\alpha-c-1)}{\Gamma(n\lambda+\alpha)} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right)^{c+1} \quad (33)$$

6. BAYES ESTIMATORS UNDER $g_2(\theta)$

From equation (6), on using (30), the Bayes estimator of θ under squared error loss function is given by

$$\hat{\theta}_s = \frac{1}{(n\lambda+\alpha-1)} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right) \quad (34)$$

From equation (8), on using (31), the Bayes estimator of θ under precautionary loss function is given by

$$\hat{\theta}_p = [(n\lambda+\alpha-1)(n\lambda+\alpha-2)]^{-\frac{1}{2}} \left(\beta + \sum_{i=1}^n \log\left(\frac{x_i}{c}\right)\right) \quad (35)$$

From equation (10), on using (32), the Bayes estimator of θ under entropy loss function is given by

$$\hat{\theta}_E = \frac{1}{(n\lambda + \alpha)} \left(\beta + \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \right) \quad (36)$$

From equation (12), on using (30) and (32), the Bayes estimator of θ under K-loss function is given by

$$\hat{\theta}_K = \left[(n\lambda + \alpha - 1)(n\lambda + \alpha) \right]^{-\frac{1}{2}} \left(\beta + \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \right) \quad (37)$$

From equation (14), on using (29) and (33), the Bayes estimator of θ under Al-Bayyati's loss function is given by

$$\hat{\theta}_{Al} = \frac{1}{(n + \alpha - c - 1)} \left(\beta + \sum_{i=1}^n \log \left(\frac{x_i}{c} \right) \right) \quad (38)$$

CONCLUSION

In this paper, we have obtained a number of estimators of parameter of Gamma-Pareto distribution. In equation (4) we have obtained the maximum likelihood estimator of the parameter. In equation (23), (24), (25), (26) and (27) we have obtained the Bayes estimators under different loss functions using quasi prior. In equation (34), (35), (36), (37) and (38) we have obtained the Bayes estimators under different loss functions using gamma prior. In the above equation, it is clear that the Bayes estimators depend upon the parameters of the prior distribution. We therefore recommend that the estimator's choice lies according to the value of the prior distribution which in turn depends on the situation at hand.

References

- [1] Alzaatreh, A., Famoye, F. and Lee, C., (2012). "Gamma-Pareto distribution and its Applications". *Journal of Modern Applied Statistical Methods*, Vol. 11, No. 1, 78-94.
- [2] Zellner, A., (1986). "Bayesian estimation and prediction using asymmetric loss functions". *Jour. Amer. Stat. Assoc.*, 91, 446-451.
- [3] Basu, A. P. and Ebrahimi, N., (1991). "Bayesian approach to life testing and reliability estimation using asymmetric loss function". *Jour. Stat. Plann. Infer.*, 29, 21-31.
- [4] Norstrom, J. G., (1996). "The use of precautionary loss functions in Risk Analysis". *IEEE Trans. Reliab.*, 45(3), 400-403.

-
- [5] Calabria, R., and Pulcini, G. (1994). "Point estimation under asymmetric loss functions for left truncated exponential samples". *Comm. Statist. Theory & Methods*, 25 (3), 585-600.
 - [6] D.K. Dey, M. Ghosh and C. Srinivasan (1987). "Simultaneous estimation of parameters under entropy loss". *Jour. Statist. Plan. and infer.*, 347-363.
 - [7] D.K. Dey, and Pei-San Liao Liu (1992): "On comparison of estimators in a generalized life Model". *Microelectron. Reliab.* 32 (1/2), 207-221.
 - [8] Wasan, M.T., (1970): "Parametric Estimation". New York: Mcgraw-Hill.
 - [9] Al-Bayyati, (2002): "Comparing methods of estimating Weibull failure models using simulation". Ph.D. Thesis, College of Administration and Economics, Baghdad University, Iraq.