

Occurrence and Pathogenicity of Fungal Pathogens in Guinea Corn, Millet and Wheat Seeds and their Control with Extracts of Garlic (*Allium sativum*) Cloves and Pawpaw (*Carica papaya*) Seeds

GWA, V. I.^{AB,*} AND WILLIAMS, E. E.^B

^aDepartment of Agronomy, Faculty of Agriculture, Federal University of Lafia, PMB 146, Lafia, Nasarawa State, Nigeria

^bDepartment of Crop Protection, Faculty of Agriculture, Federal University Dutsin-Ma, PMB 5001, Katsina State, Nigeria

*E-mail: gwavictor9@gmail.com

Abstract: This study investigated the occurrence, pathogenicity, and *in vitro* control of fungal pathogens associated with wheat, sorghum, and millet seeds using garlic (*Allium sativum*) cloves and pawpaw (*Carica papaya*) seed extracts. Extracts were tested at concentrations of 0g/200 mL, 50g/200 mL, 100g/200 mL, and 150g/200 mL, with experiments conducted in a completely randomized design, replicated three times, and incubated at $28 \pm 2^\circ\text{C}$ for five days. The fungal pathogens identified included *Aspergillus niger*, *A. flavus*, and *Lasiodiplodia theobromae*, with *A. niger* showing the highest frequency in sorghum (53.33%), millet (58.33%), and wheat (68.75%), while *L. theobromae* had the lowest frequency in these cereals (sorghum: 26.67%, millet: 16.67%, wheat: 12.50%). Pathogenicity tests revealed that *A. niger* and *L. theobromae* were highly pathogenic in wheat and very highly pathogenic in sorghum and millet. Garlic and pawpaw extracts effectively inhibited the radial growth of these fungi, with the highest concentration (150g/200 mL) exhibiting the greatest antifungal activity. Significant differences ($P \leq 0.05$) were observed for growth inhibition across incubation days, extract types, and fungal pathogens. The findings suggest that garlic and pawpaw extracts at optimized concentrations can serve as eco-friendly alternatives for managing fungal pathogens in cereals, particularly wheat, sorghum, and millet. Formulating these extracts into sustainable seed treatment technologies could help mitigate post-harvest losses, enhance food security, and align with environmentally sustainable agricultural practices

Keywords: Food security; Garlic Pathogenicity test; Pawpaw; Post-harvest

INTRODUCTION

Cereal crops such as guinea corn (*Sorghum bicolor*), millet (*Pennisetum glaucum*), and wheat (*Triticum aestivum*) are staple food sources globally, particularly in sub-Saharan Africa, where they contribute significantly to food security and livelihoods. Staple cereals such as guinea corn (*Sorghum bicolor*), millet (*Pennisetum glaucum*), and wheat (*Triticum aestivum*) constitute a significant portion of the diet in Nigeria and serve as important sources of income for millions of rural farmers. Despite

their importance, fungal infections during cultivation, storage, and processing threaten the quality and yield of these crops. This challenge is exacerbated in regions like Dutsin-Ma, Katsina State, where the climatic conditions favor fungal proliferation, necessitating urgent and sustainable intervention strategies. Pathogenic fungi, including species of *Fusarium*, *Aspergillus*, *Penicillium*, and *Colletotrichum*, have been identified as major contributors to seedborne diseases

which not only compromise seed viability but also, leading to reduced germination rates, nutritional degradation, and the production of harmful mycotoxins (Amadioha & Adeoti, 2020; Udomkun et al., 2021). Fungal pathogens are among the most destructive agents in cereals, causing significant pre- and post-harvest losses. Effective management of these pathogens is thus critical for improving crop productivity, ensuring food safety, and protecting public health.

Traditional control measures often involve chemical fungicides, which, while effective, pose environmental and health risks, including residual toxicity and the emergence of resistant fungal strains and high cost (Gwa & Akombo, 2016; Kumar et al., 2020). Consequently, there is an increasing demand for eco-friendly and sustainable alternatives, such as plant-based biofungicides (Gwa, 2021; Gwa & Iliyasu, 2024). Garlic (*Allium sativum*) and pawpaw (*Carica papaya*) are well-documented for their antifungal properties, attributed to bioactive compounds like allicin in garlic and papain in pawpaw (Lahlou et al., 2019; Mekonnen et al., 2023). The use of plant-based biofungicides, derived from natural extracts with antifungal properties, has gained increasing attention as a viable alternative to synthetic chemicals (Gwa et al., 2021). Plant such as Garlic (*Allium sativum*) and pawpaw (*Carica papaya*) are two plants with well-documented antimicrobial activities that offer promising potential for controlling fungal pathogens. Garlic contains bioactive compounds such as allicin, ajoene, and sulfur compounds, which exhibit broad-spectrum antifungal activity (Mekonnen et al., 2023). Similarly, pawpaw seeds are rich in phenolic compounds, flavonoids, and proteolytic enzymes, which have shown efficacy against a range of fungal species (Lahlou et al., 2019). The integration of these plant extracts into seed treatment protocols represents a sustainable and eco-friendly approach to disease management in cereal crops

This study investigates the occurrence, pathogenicity, and control of fungal pathogens affecting guinea corn, millet, and wheat seeds in Dutsin-Ma, Nigeria. The objectives include identifying the prevalent fungal pathogens affecting these cereals, evaluating their pathogenic potential, and assessing the antifungal efficacy of garlic and pawpaw seed extracts. Furthermore, the research seeks to contribute to sustainable agricultural practices, addressing the challenges of food security and post-harvest losses in rural communities to the development of sustainable seed treatment technologies that align with local agricultural practices and environmental conservation goals.

MATERIALS AND METHODS

Study Area

The study was conducted in Federal University Dutsin-Ma, Katsina State, Nigeria. The study area is characterized by a semi-arid climate with temperatures ranging from 25°C to 40°C, an average annual rainfall of 600–800 mm, and relative humidity conducive to fungal growth. The selection of Dutsin-Ma was based on its agricultural significance and reported challenges with diseases and cereal storage.

Collection of diseased seed samples

Guinea corn (*Sorghum bicolor*), millet (*Pennisetum glaucum*), and wheat (*Triticum aestivum*) seeds were collected from farms and stores in Dutsin-Ma. Each sample was collected in sterile polythene bags to prevent contamination. Samples were taken to Laboratory for isolation and identification of pathogens.

Preparation of Potato Dextrose Agar (PDA) medium

Potato Dextrose Agar (PDA) was prepared by dissolving 3.8 g of powdered PDA in 120 mL sterile water in a conical flask, then heated on hot plate and stirred to dissolve. The conical

flask was sealed with cotton wool and foil paper. The PDA solution was then autoclaved at 121°C for 15 minutes and allowed to cool down to 45-50°C in water bath. Exactly 0.12 g of Streptomycin Sulphate powder was added into the medium to suppress the growth of bacterial. Exactly 20 mL of the molten PDA was poured into sterile Petri dishes and allowed to solidify before inoculation of the prepared plant tissues.

Sterilization and inoculation of seed samples

Seed samples of guinea corn, millet, and wheat were surface-sterilized using 5 % Sodium hypochlorite solution, rinsed in three successive changes of distilled water, and air-dried (Gwa & Lum 2024) before inoculation of the diseased tissues on PDA. Four seeds of each sample were plated on PDA medium in Petri dishes and the plates were incubated at $28 \pm 2^\circ\text{C}$ for 5 days until fungal colonies appeared.

Isolation and identification of fungi

Fungal colonies that emerged were sub-cultured on freshly prepared PDA for another 5 days to obtain pure cultures. Morphological characteristics such as colony color, texture, and sporulation and microscopic features such as conidia shape and size were used to identify fungi based on standard manuals (Barnett & Hunter, 2003).

Determination of frequency of occurrence of the pathogen

The frequencies of the isolated pathogens were determine by counting the number of times each isolate occurred out of the total number of isolates grown on PDA during the period of isolation on each crop. This was expressed as percentage of the total number of all isolates on each crop as described in the equation below.

$$\% \text{ frequency of occurrence} = \frac{a}{b} \times \frac{100}{1}$$

a = number of times each isolate was encountered on a particular crop.

b = total number of times the different isolates were encountered on a particular crop.

Pathogenicity Test

Healthy seeds of guinea corn, millet, and wheat were surface-sterilized with 5 % Sodium hypochlorite solution, then rinsed in three successive changes of sterile distilled water to remove chlorox, air-dried and inoculated with spore suspensions (1×10^6 spores/mL) of isolated fungi. The spore suspensions (1×10^6 spores/mL) was mixed with 20 mL of PDA and allowed to solidify. Seeds were incubated on the solidified PDA and covered with moist filter paper in Petri dishes for 7 days at $28 \pm 2^\circ\text{C}$. Control seed plates were treated with sterile distilled water only and observed under same condition. Appearances of symptoms were observed for the period of incubation using a scale developed for the experiment as follows:

Table 1: Scale for pathogenicity test

Grade	Scale (%)	Symptoms	Classification of isolates
0	0-20	No or little infection	No Infection
1	21-40	Presence of infection	Slight infection
2	41-60	Infection conspicuously present	Moderate infection
3	61-80	Infection present in large quantity	High infection
4	81-100	Whole seed completely infected	Very high infection

Preparation of plant extracts

Precisely 0 g, 50 g, 100 g, and 150 g of pawpaw seed and garlic clove powders were accurately weighed using an electric weighing machine. Then, 200 mL of sterile water was measured with a measuring cylinder and poured into separate conical flasks containing the respective amounts of pawpaw seed and garlic clove powders. The mixtures were thoroughly stirred and allowed to settle for 24 hours. Afterward, they were filtered to obtain extracts with concentrations of 0 g/200 mL, 50 g/200 mL, 100 g/200 mL, and 150 g/200 mL for both pawpaw seed and garlic clove extracts.

Antifungal assay

The antifungal activity of garlic and pawpaw extracts was evaluated using the poisoned food technique. Exactly 5 mL of 0 g/200 mL, 50 g/200 mL, 100 g/200 mL, and 150 g/200 mL extract concentration of garlic and pawpaw were separately amended in 15mL of PDA and left to solidify before inoculation. Fungal plugs (5 mm diameter) of each of the isolated fungus were placed at the center of each plate. Plates were incubated at $28 \pm 2^\circ\text{C}$ for 5 days. Fungal growth was measured (diameter of colonies) using a transparent ruler in both the treated and the control plates for 5 days from the point of inoculation to the distal growth margin (Gwa & Iliyasu, 2024).

Statistical analysis

Data on fungal occurrence, pathogenicity, and inhibition rates were analyzed using analysis of variance (ANOVA) at a significance level of $P < 0.05$. The means were compared using Duncan's Multiple Range Test (DMRT).

RESULTS

Identification of fungi

Fungi identified from wheat, sorghum and millet seeds were *Aspergillus niger*, *A. flavus* and *Lasidiopodia theobromae*. Cultures of these fungi grown on Potato Dextrose Agar after 5 days of incubation are presented in Figure 1-3 below.




Figure 1




Figure 2




Figure 3

Figure 1: Culture of *A. niger* grown on PDA after 5 days of inoculation; Figure 2: Culture of *A. flavus* grown on PDA after 5 days of inoculation; Figure 3: Culture of *Lasidiopodia theobromae* grown on PDA after 5 days of inoculation

Frequency of fungal isolates

Occurrence of fungi shows that in sorghum seeds, *A. niger* had the highest frequency of eight [8] while *A. flavus* was the least occurred with frequency of only three [3]. In Wheat *A. niger* was the fungus with the highest occurrence of seven [7] compared with *L. theobromae* with the least frequency of two [2]. Similarly, millet had the highest frequency of eleven [11] isolates of *A. niger* compared with only two [2] of *L. theobromae*. This result indicated that *A. niger* is the most occurred fungus in wheat, millet and sorghum in the study area as shown in Table 2.

Pathogenicity test of *L. theobromae* in sorghum, wheat and millet

Pathogenicity test of *L. theobromae* conducted on healthy sorghum, wheat and millet showed that the fungus was able to elicit infection on the healthy seeds grown on PDA amended with spore suspension of *L. theobromae*. Growth of the fungus was apparently seen on the seeds on the PDA culture plates indicated that the seeds of wheat, sorghum and millet were severely infected as seen in Figures 4-6.

Table 2: Fungi Isolated from Sorghum, Wheat and Millet and their frequencies and percentage frequencies

Crop	Fungus	Frequency	Percentage Frequency (%)
Sorghum	<i>Aspergillus niger</i>	8	53.33
	<i>A. flavus</i>	3	20.00
	<i>Lasidiopodia theobromae</i>	4	26.67
Total		15	100
Wheat	<i>A. niger</i>	7	58.33
	<i>A. flavus</i>	3	25.00
	<i>L. theobromae</i>	2	16.67
Total		12	100
Millet	<i>A. niger</i>	11	68.75
	<i>A. flavus</i>	3	18.75
	<i>L. theobromae</i>	2	12.50
Total		16	100

**Figure 4****Figure 5****Figure 6**

Figure 4: Pathogenicity of *L. theobromae* on sorghum grown on PDA; Figure 5: Pathogenicity of *L. theobromae* on wheat grown on PDA; Figure 6: Pathogenicity of *L. theobromae* on millet grown on PDA

The levels of infectivity and growth ranged from highly pathogenic in wheat to very highly pathogenic in sorghum and millet, respectively based on the result of pathogenicity obtained and compared to the scale developed to classify the pathogenicity of the fungus identified as shown in Table 3.

Pathogenicity test of *A. niger* in sorghum, wheat and millet

Pathogenicity test of *A. niger* inoculated into healthy sorghum, wheat and millet seeds show that the fungus was able to cause infection within the 5 days of incubation on PDA.

Table 3: Pathogenicity test of *L. theobromae* inoculated into healthy sorghum, millet and wheat seeds after 5 days on PDA

Seeds	Grade	Scale	Symptoms interpretation	Classification of isolates
Wheat	3	61-80	Large quantity of seeds were infected by pathogens	Highly pathogenic
Sorghum	4	81-100	Whole seeds completely infected by pathogen	Very highly pathogenic
Millet	4	81-100	Whole seeds completely infected by pathogen	Very highly pathogenic

Symptoms of infections were observed on all the seeds with level of infectivity ranging from one crop seed to another as presented in Figures 7-9.



Figure 7



Figure 8




Figure 9

Figure 7: Pathogenicity of *A. niger* on sorghum grown on PDA; Figure 8: Pathogenicity of *A. niger* on wheat grown on PDA; Figure 9: Pathogenicity of *A. niger* on millet grown on PDA

Growth of the fungus and severity of infectivity ranged from highly pathogenic in wheat to very highly pathogenic in sorghum and millet, respectively based on the scale used to classify the virulence of pathogenicity of the fungus according to the pathogenicity test result as presented in Table 4.

Effect of concentration of garlic extract on growth inhibition of *A. niger*

The result presented in Table 5 shows that growth of *A. niger* decreased significantly as

the concentration of garlic extract increased. At 150 g/200 mL, the fungus exhibited the lowest growth across all days compared to the control (0.00 g/200 mL) and to the other concentrations.

Effect of garlic extract on growth inhibition of *L. theobromae*

Table 6 shows that the growth of *L. theobromae* was significantly reduced with increasing concentrations of garlic extract. However, radial growth was increased as the number of days for incubation increased. The highest

Table 4: Pathogenicity test of *A. niger* inoculated into healthy sorghum, millet and wheat seeds after 5 days on PDA

Seeds	Grade	Scale	Symptoms interpretation	Classification of isolates
Wheat	3	61-80	Large quantity of seed were infected by pathogens	Highly pathogenic
Sorghum	4	81-100	Whole seeds completely infected by pathogen	Very highly pathogenic
Millet	4	81-100	Whole seeds completely infected by pathogen	Very highly pathogenic

Table 5: Effect of garlic extract on *A. niger* grown on PDA 5 days of inoculation

Conc (g/200 mL)	Days after inoculation and radial growth (mm)				
		1	2	3	4 5
0.00	5.0000 ^a	18.0000 ^a	31.0000 ^a	40.0000 ^a	52.0000 ^a
50.00	1.5000 ^{cd}	11.0000 ^b	21.0000 ^b	31.0000 ^a	42.5000 ^a
100.00	2.0000 ^{bc}	10.0000 ^b	13.0000 ^c	18.0000 ^b	26.5000 ^b
150.00	3.0000 ^b	9.0000 ^b	15.8333 ^c	15.5000 ^b	22.5000 ^b
SEM	0.4269	1.1415	2.2491	3.0252	3.6323

Values followed by the same letter in each column indicate no significant difference (using Duncan’s Multiple Range Test, DMRT).

Table 6: Effect of garlic extract on *L.theobromae* growth on PDA after 5 days of inoculation

Conc (g/200 mL)	Days after inoculation and radial growth (mm)				
	1	2	3	4	5
0.00	5.0000 ^a	18.0000 ^a	31.0000 ^a	40.0000 ^a	51.5000 ^a
50.00	2.5000 ^b	3.0000 ^b	4.5000 ^b	5.5000 ^b	5.5000 ^b
100.00	1.5000 ^{bc}	2.0000 ^b	2.5000 ^b	2.5000 ^b	2.5000 ^b
150.00	1.0000 ^c	1.5000 ^b	2.0000 ^b	2.0000 ^b	2.0000 ^b
SEM	0.4885	2.1071	3.7004	4.8112	6.3048

Values followed by the same letter in each column indicate no significant difference (using Duncan's Multiple Range Test, DMRT).

growth was observed in the control (0.00 g/200 mL), while the least growth occurred at the highest garlic concentration (150 g/200 mL).

Effect of Pawpaw Seed extract on growth inhibition of *A niger*

Result presented in Table 7 shows increasing

concentrations of pawpaw seed extract reduced the growth of *A. niger*. The control (0.00 g/200 mL) recorded the highest fungal growth, while the lowest growth was observed with 150 g/200 mL of extract, especially on Day 5. Significant differences among concentrations were observed on each day of incubation

Table 7: Effect of pawpaw seed extract on *A. niger* growth on PDA after 5 days of inoculation

Conc (g/200 mL)	Days after inoculation and radial growth (mm)				
	1	2	3	4	5
0.00	5.0000 ^a	18.0000 ^a	31.0000 ^a	40.0000 ^a	51.5000 ^a
50.00	3.0000 ^b	10.0000 ^b	22.0000 ^b	30.0000 ^b	42.5000 ^a
100.00	2.0000 ^b	13.5000 ^b	20.0000 ^b	22.5000 ^c	26.5000 ^b
150.00	3.0000 ^b	12.0000 ^b	17.0000 ^b	20.5000 ^c	23.5000 ^b
SEM	0.3718	0.9791	1.6719	2.3415	3.4826

Values followed by the same letter in each column indicate no significant difference (using Duncan's Multiple Range Test, DMRT).

Effect of Pawpaw Seed extract on growth inhibition of *L. theobromae*

Result in Table 8 shows that the growth of *L.theobromae* decreased significantly with increasing concentrations of pawpaw

seed extract. The control (0.00 g/200 mL) consistently recorded the highest fungal growth, while the lowest growth occurred at 150 g/200 mL of extract, especially on Day 5. Significant differences were observed among concentrations on each day of incubation

Table 8: Effect of pawpaw seed extract on *L. theobromae* growth on PDA after 5 days of inoculation

Conc (g/200 mL)	Days after inoculation and radial growth (mm)				
	1	2	3	4	5
0.00	5.0000 ^a	18.0000 ^a	31.0000 ^a	40.0000 ^a	51.5000 ^a
50.00	1.0000 ^b	10.0000 ^b	20.5000 ^b	31.5000 ^a	45.0000 ^a
100.00	.0000 ^b	3.5000 ^c	5.5000 ^c	8.5000 ^b	18.0000 ^b
150.00	.0000 ^b	2.0000 ^c	5.0000 ^c	9.0000 ^b	13.0000 ^b
SEM	0.6336	1.9415	3.3274	4.1939	5.0370

Values followed by the same letter in each column indicate no significant difference (using Duncan's Multiple Range Test, DMRT).

DISCUSSION

The results of the findings revealed that *A. niger*, *A. flavus* and *L. theobromae* are the

cause of infection in major cereal crops like guinea corn, millet, and wheat in the study area. These pathogens significantly affect the

productivity and quality of the crops both in the field and in storage. This result agreed with findings obtained by Kumar et al. 2019 who isolated fungal species like *Fusarium spp.*, *Aspergillus spp.*, *Penicillium spp.*, and *Alternaria spp.* in these crops. According to Adebayo & Omotayo (2021) the fungi are not only capable of reducing seed viability but also have the ability to pose health challenges as a result of secretion of toxic substances like mycotoxins, aflatoxins and fumonisins. These fungi pathogens are commonly found in semi-arid regions like Dutsin-Ma, Nigeria, even on leguminous crops like cowpea (Mamkaa & Gwa, 2018) and on vegetables like tomato and pepper (Sani & Gwa, 2018; Zubairu & Gwa, 2019). The high frequency of fungal occurrence observed in the study area may be exacerbated by suboptimal post-harvest storage practices by peasant farmers and in some instances due to high humidity levels (Usman et al., 2022).

Pathogenicity tests of *A. niger* and *L. theobromae* conducted using wheat, sorghum and millet showed that the fungi elicited infections in all the crop seeds with the highest infection in sorghum and millet compared to wheat. This may be due to the high aggressive and virulent nature of the pathogens on the substrate resulting to more infection (Gwa & Okrikata, 2019). It has been found on millet and wheat that infections by these pathogens have resulted in reduced seed vigour and germination rate by 40-60 %. (Mukhtar et al., 2020).

The study revealed that period of incubation and concentration of extracts from *Allium sativum* (garlic) and *Carica papaya* (pawpaw) have significantly good result due to their antifungal properties. Results have shown that garlic contained bioactive compounds such as allicin, ajoene, and diallyl disulfide, which disrupt fungal cell membranes and inhibit spore germination (Iram et al., 2021). Similarly, finding by Bala et al. (2022) showed that a 10 % aqueous extract

of garlic inhibited the growth of *A. flavus* and *Fusarium oxysporum* in wheat by over 80 %. The effectiveness of pawpaw seeds on the other hand depends on the presence phytochemical compounds alkaloid, flavonoid, papain which contain antifungal properties. Research by Omodara et al. (2023) showed that as the concentrations of pawpaw seed extracts increased, fungal growth in millet and guinea corn reduced by 70 % yielding better results. It has further being demonstrated that the combination of garlic and pawpaw extracts produced a synergistic effect, enhancing antifungal efficacy of the extracts. According to, Aliyu et al. (2023) the combination of 5 % garlic extract with 5 % pawpaw seed extract reduced the pathogenicity of *Fusarium spp.* in millet by 85 %. Such combinations could be effective in the management of mixed fungal infection observed by a farmer.

It is therefore critical in reducing fungal infections in cereal crops which will ultimately improve food security in Nigeria. Appropriate and effective use of botanical extracts of garlic and pawpaw can reduce post-harvest losses associated with pathogenic attack, enhance seed viability for a long period of time, and reduce over dependence on synthetically hazardous chemicals, which are often costly and non-biodegradable in the environment

CONCLUSION

The study has shown that *Aspergillus niger*, *A. flavus*, *Lasiodiplodia theobromae* are pathogens of sorghum, millet and wheat. The result further revealed that *A. niger* occurred more in all the crops compared to the other pathogens. The occurrence and pathogenicity of fungal pathogens in guinea corn, millet, and wheat seeds present significant challenges to food security in Dutsin-Ma, Nigeria. However, botanical extracts from garlic and pawpaw offer a promising, eco-friendly solution for fungal control. Use of these extracts could provide enhanced antifungal effects, reduce

economic losses, and promote sustainable agricultural practices in the region.

CONFLICT OF INTEREST DISCLOSURE

The authors of this research article declare that there were no conflicts of interest.

FUNDING ACKNOWLEDGEMENT

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

REFERENCES

- Adebayo, T., & Omotayo, O. (2021). Mycotoxins and their impact on food security in Nigeria. *Journal of Agricultural Research*, 8(4), 45-56.
- Akpan, E. O., & Idowu, M. O. (2024). Garlic and pawpaw as biofungicides in cereal farming: A case study in Nigeria. *Journal of Agricultural Science and Technology*, 10(1), 101-108.
- Aliyu, H. A., et al. (2023). Synergistic effects of garlic and pawpaw seed extracts on fungal pathogens of cereal crops. *African Journal of Biotechnology*, 22(3), 78-86.
- Amadioha, A. C., & Adeoti, E. I. (2020). Impact of fungal diseases on cereal crop yields in sub-Saharan Africa. *African Journal of Agricultural Research*, 15(5), 341-349.
- Bala, K., et al. (2022). Antifungal properties of garlic extract against wheat pathogens. *Plant Pathology Research*, 10(2), 123-129.
- Duru, C. C., & Nwosu, J. I. (2019). Biocontrol of fungal pathogens in cereals using indigenous plants. *African Crop Science Journal*, 27(4), 421-432.
- Eze, O., & Ukwuru, M. (2020). Advances in biofungicides for sustainable agriculture. *Nigerian Agricultural Journal*, 15(3), 89-98.
- Eze, P. C., & Onu, E. N. (2023). Sustainable control of post-harvest diseases in cereals. *Journal of Agricultural Innovation*, 8(7), 215-223.
- Gwa, V. I., & Okrikata, E. (2019). Occurrence of fungal pathogens of Ogoja and Pepa white yam (*Dioscorea rotundata* Poir.) tuber cultivars in Zaki-Biam, Nigeria. *Nigerian Journal of Plant Protection*, 33(2), 102-112.
- Gwa, V. I. (2021). Botanicals in plant disease control. Chapter 3, pages 53-82. In: T.O. Adejumo and R.F. Voegelé (Eds): *Biopesticides: Botanicals and Microorganisms for Improving Agriculture and Human Health*. Logos Verlag, Berlin, Germany. 294pp. ISBN: 978-3-8325-5264-0.
- Gwa, V. I., Muhammad Akram, & Naveed Munir (2021). Effect of selected medicinal plant extracts and synthetic antifungal agents on post-harvest rot fungi of Hembankwase white yam (*Dioscorea rotundata* Poir) tuber cultivar. *Nigerian Journal of Mycology*, 13, 1-11.
- Gwa, V. I., & Iliyasu, A. (2024). Potency of aloe vera (*Aloe barbadensis* Miller) and bell pepper (*Capsicum annuum* L.) in the control of *Aspergillus niger* causing black mold disease of processed groundnut (*Arachis hypogaea* L.) seeds. *Plant Pathology & Quarantine*, 14(1), 60-70. <https://doi.org/10.5943/ppq/14/1/5>.
- Gwa, V. I., Lum, A. F. (2024). Bioactivity of plant extracts against *Fusarium oxysporum* f. sp. *Lycopersici* (Sacc.) causing wilt disease of tomato (*Solanum lycopersicum* L.) in the southern Guinea savannah, Nigeria. *Journal of Plant Science and Crop Protection*, 7(1), 103.
- Iram, S., et al. (2021). Mechanisms of action of garlic-derived bioactives against plant pathogens. *Frontiers in Plant Science*, 12, 2345.
- Khan, N., Ghori, S. A., & Abbas, S. (2022). Role of plant extracts in managing seedborne fungi. *Pakistan Journal of Botany*, 54(5), 1501-1510.
- Kumar, R., et al. (2019). Fungal contamination of cereals: A review. *Agricultural Reviews*, 40(4), 303-310.
- Kumar, S., Yadav, P., & Gupta, J. (2020). Challenges in controlling seedborne fungal pathogens of cereals. *Plant Pathology Journal*, 36(2), 99-110.
- Lahlou, F., El Bakali, A., & Bouchal, H. (2019). Natural antifungal agents from plant sources. *Current Research in Food Science*, 3(1), 12-24.
- Mamkaa, D. P., & Gwa, V. I. (2018). Effect of *Moringa oleifera* and *Vernonia amygdalina* leaf extracts against *Aspergillus flavus* and *Botryodiplodia theobromae* causing rot of cowpea (*Vigna unguiculata* (L.) Walp) seeds. *Applied Science Research Review*, 5(1,2), 1-7. DOI: 10.21767/2394-9988.100067.
- Mekonnen, H. T., Desta, B., & Abera, M. (2023). Bioactive compounds in garlic and their role in fungal inhibition. *Journal of Agricultural Science*, 15(4), 189-202.
- Mukhtar, T., et al. (2020). Pathogenicity of *Fusarium* spp. on cereal seeds. *Pakistan Journal of Botany*, 52(5), 1713-1720.
- Omodara, T. A., et al. (2023). Evaluation of pawpaw seed extracts as biocontrol agents for

- millet pathogens. *International Journal of Tropical Agriculture*, 41(1), 55-65.
- Oluwafemi, A. T., & Adeyanju, S. O. (2022). Evaluating the antifungal efficacy of pawpaw seed extracts in crop protection. *Nigerian Journal of Plant Protection*, 36(3), 44-52.
- Sani, S., & Gwa, V. I. (2018). Fungicidal effect of *Azadirachta indica* and *Zingiber officinale* extracts in the control of *Fusarium oxysporum* and *Rhizoctonia solani* on tomato (*Solanum lycopersicum*) fruits. *Innovative Techniques in Agriculture*, 2(4), 439-448.
- Usman, Y., et al. (2022). Environmental factors influencing fungal infections in stored cereals in Nigeria. *Journal of Food Safety*, 25(6), 322-334.
- Udomkun, P., Wiredu, A. N., & Nagle, M. (2021). Mycotoxin contamination in cereals: Implications for food safety in developing countries. *Food Control*, 121, 107585.
- Yusuf, A. A., & Mohammed, Z. A. (2023). Post-harvest fungal pathogens in cereal grains: An overview. *International Journal of Food Microbiology*, 169, 68-76.
- Zhang, W., Liang, C., & Yang, X. (2021). Eco-friendly fungicides in sustainable agriculture. *Agricultural Research*, 10(2), 123-136.
- Zubairu, T., & Gwa, V. I. (2019). Antifungal activity of *Azadirachta indica* A. Juss and *Moringa oleifera* L. seed extracts against rot fungi of hot pepper (*Capsicum annuum* L.) fruits in Dutsin-Ma, Katsina State, Nigeria. *FUDMA Journal of Agriculture and Agricultural Technology*, 5(2), 254-265.